

Tandem duplications and the limits of natural selection in *Drosophila yakuba* and *Drosophila simulans*

Rebekah L. Rogers¹, Julie M. Cridland², Ling Shao³, Tina T. Hu⁴,
Peter Andolfatto⁴, and Kevin R. Thornton³

Research Article

Biological Sciences, Population Biology

- 1) Ecology and Evolutionary Biology, University of California, Berkeley
- 2) Ecology and Evolutionary Biology, University of California, Davis
- 3) Ecology and Evolutionary Biology, University of California, Irvine
- 4) Ecology and Evolutionary Biology and the Lewis Sigler Institute for Integrative Genomics, Princeton University

Running head: Tandem duplications in non-model *Drosophila*

Key words: Gene duplications, *Drosophila yakuba*, *Drosophila simulans*, evolutionary novelty, population genomics, parallel adaptation, convergent evolution, mutation limited evolution, rapid evolution, Red Queen dynamics

Corresponding author: Rebekah L. Rogers, Dept. of Ecology and Evolutionary Biology, 5323 McGaugh Hall, University of California, Irvine, CA 92697

Phone: 949-824-0614

Fax: 949-824-2181

Email: rogersrl@uci.edu

Abstract

Tandem duplications are an essential source of genetic novelty, and their variation in natural populations is expected to influence adaptive walks. Here, we describe evolutionary impacts of recently-derived, segregating tandem duplications in *Drosophila yakuba* and *Drosophila simulans*. We observe an excess of duplicated genes involved in defense against pathogens, insecticide resistance, chorion development, cuticular peptides, and lipases or endopeptidases associated with the accessory glands, suggesting that duplications function in Red Queen dynamics and rapid evolution. We document evidence of widespread selection on the *D. simulans* X, suggesting adaptation through duplication is common on the X. Despite the evidence for positive selection, duplicates display an excess of low frequency variants consistent with largely detrimental impacts, limiting the variation that can effectively facilitate adaptation. Although we observe hundreds of gene duplications, we show that segregating variation is insufficient to provide duplicate copies of the entire genome, and the number of duplications in the population spans 13.4% of major chromosome arms in *D. yakuba* and 9.7% in *D. simulans*. Whole gene duplication rates are low at 1.17×10^{-9} per gene per generation in *D. yakuba* and 6.03×10^{-10} per gene per generation in *D. simulans*, suggesting long wait times for new mutations on the order of thousands of years for the establishment of sweeps. Hence, in cases where adaption depends on individual tandem duplications, evolution will be severely limited by mutation. We observe low levels of parallel recruitment of the same duplicated gene in different species, suggesting that the span of standing variation will define evolutionary outcomes in spite of convergence across gene ontologies consistent with rapidly evolving phenotypes.

Introduction

Tandem duplications are an essential source of genetic novelty that is useful for the development of novel traits (Conant and Wolfe 2008, Ohno 1970) and their prevalence in populations is therefore expected to influence the arc of evolutionary trajectories. The observed landscape of tandem duplications in *Drosophila* spans only a few percent of the genome (Rogers et al. 2014, Emerson et al. 2008, Cardoso-Moreira et al. 2011, Dopman and Hartl 2007), and it is unclear to what extent duplications, whether from newly arising or from standing variation, can provide a sufficient source of adaptive genetic variation. Tandem duplications produce a variety of novel gene structures including chimeric genes, recruited non-coding sequence, dual promoter genes, and whole gene duplications (Rogers et al. 2014, Zhou et al. 2008, Katju and Lynch 2006). Surveys within single genomes have suggested that whole gene duplications may form at low rates in comparison with SNPs, with even lower mutation rates for complex variants such as chimeric genes (Rogers, Bedford and Hartl 2009, Rogers et al. 2014, Zhou et al. 2008). Yet, these alternative genetic structures are known forces of evolutionary innovation. Whole gene duplications often develop novel functions or specialize in ancestral functions (Conant and Wolfe 2008), and chimeric genes

are more likely still to produce novel molecular effects and play a role in adaptive evolution (Rogers and Hartl 2012). Although these variants contribute substantially to the evolution of genome content (Zhou et al. 2008, Rogers, Bedford and Hartl 2009, Lynch and Conery 2000), their lower rates of formation may render evolution of tandem duplications more likely to be limited by mutation.

If population-level mutation rates are sufficiently large, new mutations will accumulate quickly and adaptation is expected to proceed rapidly (Hermisson and Pennings 2005). However, if population-level mutation rates are low, then there will be long wait times until the next new mutation and evolutionary trajectories are likely to stall at suboptimal solutions during the mutational lag (Maynard Smith 1971, Gillespie 1991, Hermisson and Pennings 2005). *Drosophila* have large population sizes in comparison to other multicellular eukaryotes with $N_e \approx 10^5 - 10^6$ (Kreitman 1983, Bachtrog et al. 2006, Andolfatto, Wong and Bachtrog 2011a) and absolute numbers of individuals large enough to provide large numbers of SNPs at many sites every generation (Karasov, Messer and Petrov 2010). However, the prevalence of other types of mutations beyond SNPs has not been systematically surveyed. If the supply of tandem duplications is limited by mutation, we expect to see suboptimal outcomes in adaptive walks, limited ability to adapt to changing environments, and low rates of evolution through parallel recruitment of the same genetic solutions in different species. The *Drosophila* offer an excellent model system for population genomics, allowing for a whole genome survey of the genetic landscape of standing variation across species in natural populations and determination of genetic convergence across taxa. There are

multiple sequenced reference genomes for *Drosophila*, and genomes are small and compact, allowing for whole genome population surveys using next generation sequencing. Here, we focus on *D. yakuba* and *D. simulans*, which are separated by 12 MY of divergence (Tamura, Subramanian and Kumar 2004), allowing for surveys of distantly related groups which are not expected to share polymorphic variation due to ancestry. Thus, we can measure the limits of standing variation and the incidence of parallel duplication across species, which should be broadly applicable to multicellular eukaryotic evolution.

Convergent evolution is regarded as the ultimate signal of natural selection: if the same solution is favored for a given environment then selection should result in similar phenotypes (Gould and Lewontin 1979). There are many known cases of convergent phenotypic evolution, but the understanding of convergence at the genetic level is limited to a small number of case studies across diverse clades (Stern 2013). These case studies have revealed convergent evolution through different genetic solutions in vertebrates (Chen, DeVries and Cheng 1997, Shapiro et al. 2009, Brodie 2010), and arthropods (Khadjeh et al. 2012, Wittkopp et al. 2003, Tanaka, Barmina and Kopp 2009, Zhen et al. 2012). Parallel evolution through similar genetic solutions, however, appears to be more common at mutational hotspots where high mutation rates at targeted sites produce mutations at a steady rate (Riehle, Bennett and Long 2001, Chan et al. 2010, Moxon, Lenski and Rainey 1998). Beyond these results from natural populations, convergence has often been observed in experimental evolution and is considered a signal of selection favoring alleles (Riehle, Bennett and Long 2001, Moxon, Lenski and Rainey 1998, Burke et al. 2010, Woods et al.

2006, Tenaillon et al. 2012). However, most studies of laboratory evolution take advantage of microbes or viruses with large population sizes roughly $10^9 - 10^{10}$ such that every mutation is likely to be sampled every generation (Riehle, Bennett and Long 2001, Moxon, Lenski and Rainey 1998) or from small populations that share a common pool of standing variation (Burke et al. 2010, Orozco-terWengel et al. 2012) and may therefore be qualitatively different outcomes in comparison to natural evolution in multicellular eukaryotes. Indeed, known examples of evolution through parallel recruitment of the same genetic solutions in natural populations often occur through a common ancestral genetic pool (Colosimo et al. 2005) or through introgression (Martin et al. 2012).

The instance of convergent evolution across unrelated taxa that do not share ancestry is essential to understanding the ways mutation limits evolution, the role of standing variation in evolutionary trajectories, and the genetic architecture of adaptation. Here, we survey standing variation for tandem duplications in *Drosophila yakuba* and *Drosophila simulans* and the role that this standing variation plays in adaptive evolution in natural populations. We identify an overrepresentation of tandem duplications with gene ontologies consistent with rapidly evolving phenotypes, signals of reduced diversity surrounding tandem duplications, and an overabundance of high frequency variants on the *D. simulans* X chromosome, pointing to a role for adaptation through gene duplication. We further show that the span of tandem duplications in populations is limited to a small fraction of the genome and that low mutation rates will lead to long wait times for sweeps on new mutations. These results imply that evolution by tandem duplication will be limited by mutation and

that parallel recruitment of gene duplicates across species is likely to be exceedingly rare even in the face of strong selection on similar phenotypes in different species.

Results

We previously identified hundreds to thousands of segregating duplications in natural populations of *D. yakuba* and *D. simulans*, including large numbers of gene duplications (Table S1) (Rogers et al. 2014). We assess the numbers and types of gene duplications, differences in duplication rates across species and explore the limits of the landscape of standing variation for tandem duplications present in each species to determine the extent to which these variants can serve as a source of genetic novelty.

Recently derived, segregating tandem duplications were detected using paired end reads and coverage changes in *D. yakuba* and *D. simulans* in samples of 20 isofemale lines derived from natural populations of each species (Rogers et al. 2014). Using divergently oriented paired-end reads, we identified 1415 segregating tandem duplications in *D. yakuba*, in comparison to 975 in *D. simulans*. The tandem duplications identified across these sample strains cover 2.6% of assayable the genome of the X and 4 major autosomes in *D. yakuba* and 1.8% of the assayable genome of the X and 4 major autosomes in *D. simulans*. These variants have a high (96%) confirmation rate (Rogers et al. 2014) and constitute a high quality dataset for population genomics.

Widespread selection on the *D. simulans* X chromosome

If tandem duplications are common targets for adaptation and selective sweeps, we should observe a shift in the site frequency spectrum (SFS) toward high frequency variants relative to neutral markers (Nielsen et al. 2005). We compare the SFS for duplications with the SFS for SNPs from 8-30 bp of short introns used as a putatively neutral proxy to determine whether duplicates are subject to selection (Figure S1). The SFS for duplications is significantly different from that of intronic SNPs on the *D. simulans* autosomes ($W = 268$, $P = 2.981 \times 10^{-6}$) and *D. yakuba* autosomes ($W = 212$, $P = 3.507 \times 10^{-6}$). In *D. yakuba* the SFS for duplicates on the X is significantly different from that of SNPs ($W = 211$, $P = 4.781 \times 10^{-4}$) using a Wilcoxon sign rank test. Duplicates show an excess of singleton variants on the autosomes in both species (Figure S1), suggesting deleterious impacts on average. We find a significant difference between the SFS of duplicates on the X chromosome and the autosomes in *D. yakuba* ($W = 172$, $P = 0.0128$) but not in *D. simulans* ($W = 183.5$, $P = 0.1848$) (Figure 1, Table S2-S3). In contrast, the SFS for duplicates on the *D. simulans* X chromosome exhibit fewer singletons in comparison to neutral SNPs and a peak of high frequency duplicates (Figure 1). We observe an excess of duplications at a sample frequency of 20 out of 20 sample strains (with no indication of duplication or misassembly in the resequenced reference) on the X chromosome of *D. simulans* in comparison to neutral SNPs ($P < 10^{-6}$), but observe no duplicates at a sample frequency of 20 out of 20 in *D. yakuba* on the X or autosomes. Furthermore comparisons of the SFS show an excess of highest frequency variants ≥ 16 out of 17 on the X ($\chi^2 = 21.8334$, $df = 1$, $P = 2.974 \times 10^{-6}$).

The excess of high frequency duplicates on the *D. simulans* X chromosome is indicative of selection favoring large numbers of tandem duplicates. These results imply that adaptation through duplication is common on the *D. simulans* X.

We have calculated average heterozygosity per site (θ_π) (Tajima 1983), Watterson's θ ($\frac{S}{a}$ per site) (Watterson 1975), and Tajima's D (Tajima 1989) for the four major autosomes and the X chromosome in *D. yakuba* and *D. simulans* using 5 kb windows with a 500 bp slide correcting for the number of sites with coverage sufficient to confidently identify SNPs (Figure S2-S11). We compare θ_π in regions surrounding tandem duplications and putatively neutral SNPs from 8-30 bp of short introns to search for signals of reduced diversity consistent with selection acting on tandem duplications. These tandem duplications are polymorphic and represent putative sweeps in progress and such comparisons to within-genome controls of neutral SNPs offer greater power than alternative tests of selection (Coop and Ralph 2012). We compare estimates of θ_π for windows from 50 kb upstream to 50kb downstream of a duplicated locus with comparable windows surrounding derived segregating SNPs from putatively neutral sites of short introns, excluding centromeric regions. We find a reduction in θ_π per site surrounding newly arisen duplications on *D. yakuba* chromosome 3R (single tailed Wilcox test, $P = 0.0064$, see Table S4, Figure 2) as well as on *D. simulans* chromosomes 2L, 3L, 3R, and the *D. simulans* X (single tailed Wilcox test, $P \leq 0.05$; see Table S4) (Figure 3). Chromosome 3L has reduced diversity surrounding duplicates in *D. simulans*, even though a region encompassing multiple duplications with signals of a broad selective sweep encompassing multiple loci at roughly 8.5 Mb is excluded (Figure S9). A significant

excess of diversity is seen on chromosome 2L in *D. yakuba* ($W = 31594$, $P = 2.517 \times 10^{-4}$), a possible product of alternative evolutionary dynamics driven by segregating inversions on 2L (Lemeunier and Ashburner 1976, Llopart, Lachaise and Coyne 2005). We further compare θ_π for windows centered about duplications that capture solely intergenic sequence with those that capture coding sequence but respect gene boundaries, and those that create chimeric genes or recruited non-coding sequence. In *D. simulans*, gene duplications that do not create chimeric genes have reduced diversity ($W = 17880$, $P = 0.01233$) in comparison to mutations that capture intergenic mutations (Figure 4C-D), but chimeric gene duplications do not ($W = 5020$, $P = 0.5755$), consistent with selection driving an excess of whole gene duplications in *D. simulans* (Rogers et al. 2014), but relationships in *D. yakuba* are not significant for chimeric gene mutations ($W = 2656$, $P = 0.8226$) or whole gene duplications ($W = 6387$, $P = 0.8847$). All of these mutations are polymorphic indicating sweeps in progress or neutral dynamics and power may be limited to detect differential selection on different types of mutations. Based on a binomial test, we observe a marginally significant overrepresentation of duplications that capture gene sequences vs solely non-coding sequences in *D. yakuba* ($P=0.029$) and highly significant for *D. simulans* ($P = 9.044 \times 10^{-5}$).

While demography and neutral evolutionary forces can result in shifts of diversity and site frequency spectra, these forces should affect sequences across individual chromosome arms and act similarly on intronic SNPs, and are therefore unlikely to explain the observed differences between duplicates and intronic SNPs. Therefore, it seems likely that the overabundance of high frequency variants on the *D. simulans* X is driven by natural

selection. Thus, we would expect many of these high frequency variants to be strong candidates for ongoing selective sweeps. We further observe large numbers of singleton variants among tandem duplicates in comparison with intronic SNPs in *D. yakuba* and *D. simulans* autosomes. Copy number variants are subject to purifying selection in *D. melanogaster* (Emerson et al. 2008, Cridland and Thornton 2010), and we observe large numbers of singleton variants in excess of neutral expectations, indicating negative selection preventing variants from rising to higher frequency. Hence, while some variants are likely to offer a means of adaptive change, many are likely to ultimately be lost from the pool of standing variation. We suggest that tandem duplications are likely to confer phenotypic impacts that are on average large enough to surpass the threshold of near neutral effects.

Rapid Evolution

Biases in the rates at which duplications form in different genomic regions or a greater propensity for selection to favor duplications in specific functional classes can result in a bias in gene ontology categories among duplicated genes. We use DAVID gene ontology analysis software to identify overrepresented functions among duplicate genes in *D. yakuba* and *D. simulans* (Rogers et al. 2014). While some categories are specific to each species, immune response and toxin metabolism are overrepresented in both species (Figure 5A, Table S5), functions that have also been identified as being overrepresented in a parallel study of *D. melanogaster* (Zichner et al. 2013). We also observe an overrepresentation of chitin cuticle formation and chemosensation in both *D. yakuba* and *D. simulans* (Table S5). Furthermore,

lipases and endopeptidases commonly found in the accessory glands are present among high frequency variants in both species and there are multiple independent duplications in chorion and egg development genes (Table S5), suggesting a role for duplications in sexual conflict. The overabundance of toxin metabolism genes and immune response peptides in both species as well as the overrepresentation of putative chemoreceptors, chitin based cuticle genes, endopeptidases, and oogenesis factors suggests that duplications may be key players in rapidly evolving systems. Moreover, the strong agreement in overrepresented functional categories points to strong selective pressures acting in parallel in these independently evolving species. To determine whether selection is favoring these functional classes, we identified duplications outside centromeric regions that lie in windows at or below the 5% tail of θ_π , consistent with selection reducing diversity. Among these genes in regions with reduced diversity we identify genes in both *D. yakuba* and *D. simulans* with functions in chorion or oogenesis, mating behavior, immune response and defense against bacteria, olfactory response, chitin metabolism, xenobiotics and toxin metabolism, and sperm development (Supplementary Information). The presence of genes with these functional categories is consistent with a portion of the overrepresentation in gene ontologies across all duplicates being driven at least in part by selection.

Limits of standing variation in natural populations

We observe hundreds of segregating tandem duplicates in *D. yakuba* and *D. simulans*, spanning 2.6% of assayable the genome of the X and 4 major autosomes in *D. yakuba*

and 1.8% of the assayable genome of the X and 4 major autosomes in *D. simulans*. If evolutionary trajectories depend on duplications to effect beneficial phenotypic changes, then the number of segregating variants available in the population may not contain desired variants as standing variation. We estimate the number of variants present in the entire population based on the observed sample variation in order to determine the extent to which selection will be limited by mutation. We estimate that the population contains at most 7,230 segregating tandem duplications in *D. yakuba* and 4,720 in *D. simulans* (Table S6-S7), corresponding to 13.4% of major chromosome arms in *D. yakuba* and 9.7% of major chromosome arms in *D. simulans*. Thus, the standing variation for tandem duplications will be insufficient to offer tandem duplications for every potential gene across the entire genome and for the majority of the genome ($\approx 85\%$), if a tandem duplication is required for adaptation, evolutionary trajectories must by definition rely on new mutations.

We calculate population level mutation rates θ_π ($4N_e\mu$) of 0.00277 for whole gene duplications 0.00082 for recruited non-coding sequence and 0.00088 for chimeras in *D. yakuba*. Population level mutation rates in *D. simulans* are slightly higher with 0.00291 for whole gene duplications 0.00117 for recruited non-coding sequence and 0.00041 for chimeras. In comparison, we calculate θ_π for putatively neutral intronic SNPs of 0.0138 for *D. yakuba* and 0.0280 for *D. simulans*. We use these estimates of θ to calculate the likelihood of adaptation from alleles among the standing variation rather than new mutation for a population (P_{sgv}) (Hermisson and Pennings 2005) assuming additive variants with a large selection coefficient of 1% under an additive genetic model. With such low levels of θ_π the likelihood of adaptation

from a tandem duplication among the standing variation is 2.2% in *D. yakuba* and 2.6% in *D. simulans* (Table 1), a strikingly low likelihood that standing variation offers a sufficient substrate for adaptation. Even with a massive selective coefficient of $s = 0.20$ (Karasov, Messer and Petrov 2010), the likelihood of adaptation from standing variation rather than new mutation is 3.1% for duplicates in *D. yakuba* and 3.4% in *D. simulans*. Chimeras are even more extreme with less than a 1% chance of fixation from standing variation (Table 1). In comparison, intronic SNPs have a likelihood of adaptation from standing variation of 12.1% in *D. yakuba* and 24.6% in *D. simulans* given $s=0.01$, and 15.7% in *D. yakuba* and 30.1% in *D. simulans*, given extreme selection coefficients of $s=0.20$ (Table 1). Thus, the limits of standing variation are expected to be far more severe for complex gene structures than for SNPs.

We calculate the per generation mutation rate μ per gene for whole gene duplications, considering duplicates that capture 90% or more of gene sequences, in agreement with previous methods (Rogers, Bedford and Hartl 2009). We estimate a whole gene duplication rate of 1.17×10^{-9} per gene per generation for *D. yakuba* and 6.03×10^{-10} per gene per generation for *D. simulans* (Figure 6, Table 1), in general agreement with estimates derived from surveys of duplicates in the *D. melanogaster* reference genome of 3.68×10^{-10} per gene per generation (Rogers, Bedford and Hartl 2009, Zhou et al. 2008). The rate of recruited non-coding sequence is 3.46×10^{-10} in *D. yakuba* and 2.42×10^{-10} in *D. simulans* and the rate of chimeric gene formation is equally low with 3.7×10^{-10} in *D. yakuba* and 8.52×10^{-11} in *D. simulans* (Figure 6, Table 1). We observe more duplications in *D. yakuba* in spite of

its lower N_e , yielding a duplication rate per gene in *D. yakuba* two-fold higher than that of *D. simulans*. Given these estimates of θ_π , we estimate T_e , the time to establishment of a deterministic sweep from new mutations in a population such that variants overcome the forces of drift (Gillespie 1991, Maynard Smith 1971), assuming that beneficial mutations appear at strongly selected sites at a rate equivalent to the genome-wide mutation rate. In reality not all mutations are beneficial and the rate of adaptive substitution is likely to be less common. These estimates therefore represent an lower bound on the time to adaptation through new mutation. With a small selection coefficient of $s=0.01$, in *D. yakuba* T_e would be 7270 generations (600 years at 12 generations per year) for SNPs, 36,000 generations (3000 years) for whole gene duplications, and over 100,000 generations (≥ 9500 years) for chimeric genes (Table 1). For *D. simulans*, these numbers point to a greater disparity between SNPs and duplications with T_e of 3580 generations (300 years) for SNPs, 34,400 generations (2800 years) for whole gene duplications, and 243,000 generations (20,000 years) for chimeric genes (Table 1). Under extreme selection coefficients and with the assumption that the beneficial mutation rate matches the mutation rate per site, wait times may be shorter, allowing for adaptation at SNPs in hundreds of generations (decades) and thousands of generations (centuries) for duplications (Table 1), though such extreme dynamics are unlikely to reflect the range of selection coefficients or the rate of adaptation for the genomewide (Jensen, Thornton and Andolfatto 2008, Andolfatto 2007). These estimates of mutation rates for whole gene duplications and complex gene structures point to long wait times for new mutations and a disparity in the response of duplicates and SNPs in the face of strong

selective pressures. Although the differences in mutation rates appear to be modest, they can result in additional thousands of years in the wait time for selective sweeps to establish with new mutations, resulting in limited ability to adapt to shifting selective pressures.

A Bayesian binomial estimate of the 95% lower CI suggests that for all mutations not observed across the 20 sample strains, their frequency in the population is ≤ 0.1329 , with a 50% lower CI of ≤ 0.0325 . Very rare mutations may have difficulty escaping the forces of drift in the population, especially if recessive (Haldane 1927), and therefore ultimately the number of duplications that are at frequencies high enough to establish deterministic selective sweeps will be considerably fewer than the number that exist in the population. The number of tandem duplications that have the potential to sweep to fixation may be substantially less than indicated by the number of segregating sites. Thus, the pool of standing variation in tandem duplications will provide only a limited substrate of novel genetic sequences and evolution will be limited by mutation.

Some 56 genes are partially or wholly duplicated both in *D. yakuba* and in *D. simulans*, a mere 11% of genes duplicated genes in *D. simulans* ($\frac{56}{478}$) and less than the number of genes duplicated multiple times in *D. simulans* alone, suggesting that there is little concurrence in the standing variation of the two species. That 56 genes would be shared across the two species is greater than expected given the limits of available standing variation of 478 duplicated genes in *D. simulans* and 875 in *D. yakuba* based on uniform chance ($P = 2.812 \times 10^{-8}$, binomial test) pointing to mutational or selective pressures on similar genes (SI Text). *D. simulans* shares fewer genes with *D. melanogaster* than does *D. yakuba*, a result

of fewer annotated gene models in *D. simulans* *w501* reference (Hu et al. 2012). Furthermore, a comparison to duplicate genes in *D. melanogaster* (Zichner et al. 2013) shows only 5 genes that exist among the segregating variation of tandem duplications in all three species. The mutations described here have been polarized with respect to ancestry, and are segregating meaning that they are expected to have formed very recently. As such, shared variants are the product of independent mutation in the two species, not shared ancestry. We find that 13.4% of the genome is present but unsampled in *D. yakuba* and 9.7% in *D. simulans*, indicating that the likelihood of shared, unsampled variation is low. Such unsampled alleles will be at low frequency and are unlikely to be able to establish selective sweeps. Hence, the portion of variation available for selective sweeps that is shared across species will be low, resulting in a rarity of evolution through parallel recruitment of tandem duplicates. Some genes within the genome are captured by as many as 6 independent duplications in *D. yakuba* and 32 independent duplications in *D. simulans*. There are 10 genes in *D. yakuba* and 12 genes in *D. simulans* that are captured by more than three independent duplications, and these have been excluded from mutation estimates (Table S8). These “hotspots” within the genome may have duplication rates high enough that evolution will not be subject to the same limitations with respect to standing variation and sweeps on new mutations.

Discussion

We have described the prevalence of tandem duplications in natural populations of *D. yakuba* and *D. simulans*, their frequencies in the population, and the genes that they affect. We find

that duplications show a bias towards gene ontologies associated with rapid evolutionary processes and that they commonly affect the X chromosome in *D. simulans* in comparison to the autosomes. In spite of their strong role in adaptation, we find low rates of parallel recruitment of tandem duplications across species due to low formation rates and mutation limited evolution.

Widespread positive selection on the X chromosome in *D. simulans*

We observe an excess of high frequency duplications on the *D. simulans* X chromosomes in comparison to neutral intronic SNPs as well as signs of reduced diversity surrounding duplications on the *D. simulans* X, consistent with widespread selection. Background selection (Charlesworth, Morgan and Charlesworth 1993) and hitchhiking (Smith and Haigh 1974) are not expected to act differently on duplications in comparison to SNPs and cannot explain the patterns observed. Yet, we observe significant differences between the SFS of duplicates and putatively neutral SNPs, pointing to a role for adaptation through tandem duplication. We also observe reduced nucleotide diversity surrounding tandem duplications on the *D. simulans* X, consistent with selection favoring duplicates. Hence, the overabundance of high-frequency duplications on the X is likely to be driven by selection and these represent strong candidate loci for ongoing selective sweeps. Based on the newly assembled *D. simulans* reference, X vs. autosome divergence indicates faster evolution on the X chromosome at non-synonymous sites, long introns, and UTRs (Hu et al. 2012). This pattern is distinct from observations at synonymous sites as well as general patterns of differential evolution on

the autosomes (Hu et al. 2012), further evidence of more frequent selective sweeps on the X chromosome.

The X chromosome is thought to evolve rapidly due to sexual conflict, intragenomic conflict, and sexual selection (Presgraves 2008) and thus multiple selective forces may facilitate the spread of duplicates on the X. The X chromosome in *D. simulans* houses an excess of duplicates in comparison to all autosomes, as well as a strong association with repetitive sequence and tandem duplications on the X (Rogers et al. 2014). Therefore, the X chromosome appears to be subject to particularly rapid evolution in duplicate content in *D. simulans*. Previous work has identified signals of adaptation through duplication on the *D. melanogaster* X chromosome (Thornton and Long 2002; 2005), suggesting parallel evolution through duplication in these species. However, we do not observe similar patterns in *D. yakuba*, suggesting that the X may either be evolving under different selective pressures in the two different clades or that selective pressures on the *D. yakuba* X chromosome are of lesser magnitude. Stronger sexual selection, greater selection for X-chromosome related traits, sympatric associations with competitor species with reinforcement for mating aversion, or a greater instance of driving X chromosomes might potentially drive these species differences in X-chromosome evolution, and elucidating the nature of selection on these sex chromosomes may help explain the adaptive (or selfish) role of tandem duplication on the X.

Mutation limited evolution

While both *D. simulans* and *D. yakuba* house a rich diversity of duplicated sequences, only a few percent of the genome will be covered by tandem duplications. With lower mutation rates for duplications (Lynch and Conery 2003, Rogers, Bedford and Hartl 2009, Zhou et al. 2008), there may be long wait times to achieve any single new mutation, and the landscape of standing variation will shape evolutionary outcomes. As such, any evolutionary path that is dependent upon duplications of any specific genomic sequence will be severely limited by the small likelihood that the necessary mutation is among the standing variation in tandem duplications. *Drosophila* represent organisms with large effective population sizes (Figure 6) (Bachtrog et al. 2006, Eyre-Walker et al. 2002) and are expected to host large numbers of duplications as standing variation in comparison to other multicellular eukaryotes. We have shown that the number of tandem duplications segregating in the population is substantially smaller than the number of mutations needed to guarantee a duplicate of any desired genomic region. However, when population level mutation rates are small, standing variation is unlikely to offer a sufficient substrate for selective sweeps and systems will be stuck waiting for new mutations that are slow to materialize (Hermisson and Pennings 2005). We observe population level mutation rates θ per gene for tandem duplications on the order of 0.00277 in *D. yakuba* and 0.00291 in *D. simulans* (Table 1, Figure 6) resulting in low probabilities that standing variation offers the major source of adaptation and long wait times to sweeps on new mutations on the order of hundreds to thousands of years. While retrogenes might provide additional sources of duplicated sequences, their rates of formation are exceptionally limited

(Schrider et al. 2011, Zhou et al. 2008) and they are therefore not expected to contribute more substantially than tandem duplications to genomic variation and will not suffice to overcome these limitations of low mutation rates. Thus, we conclude that outside of a small number of mutational hotspots evolution through duplication is mutation limited even in *Drosophila* which have large N_e , and that these limits are expected to be even more severe for many other multicellular eukaryotes, especially vertebrates.

The majority of tandem duplications identified in *D. yakuba* and *D. simulans* appear to be at extremely low frequency, with an excess of singleton variants in comparison to neutral intronic SNPs, suggesting that large numbers of duplications are detrimental, consistent with previous work in other species (Emerson et al. 2008). It has previously been argued that the accumulation of duplications is the product of small N_e and inability of selection to purge nearly neutral alleles from the population (Lynch and Conery 2000, Lynch 2007). However, we show that duplicates are less likely to be neutral in comparison to putatively neutral intronic SNPs suggesting that they often have phenotypic effects larger than the limit near-neutrality. We have shown that both positive and negative selection will affect the fixation or loss of duplications and that simplified nearly neutral theories are unlikely to explain the patterns observed across species. Rather, selection is expected to play an appreciable role in the evolution of tandem duplications and their contribution to genome content.

Likelihood of parallel recruitment of tandem duplications across species

Convergent evolution is often interpreted of a signal of adaptation in experimental evolution and in natural populations (Gould and Lewontin 1979, Stern 2013). Here, we show that for tandem duplications, parallel recruitment of genes for duplication and diversification independent from shared ancestry will be very rare in spite of convergence in functional categories represented. Thus, the reliance on genetic convergence to establish natural selection in natural populations will underreport selected alleles and result in significant underestimation of the number and types of alleles that are selected. Though convergence is common in experimental evolution of both prokaryotic systems and multicellular eukaryotes with shared ancestry, these results suggest that such convergence is unlikely to reflect the frequency of convergent evolution in natural populations of independently evolving species of multicellular eukaryotes that have little shared standing variation. We observe an excess of variants with gene ontologies consistent with similar rapid evolutionary processes both in *D. yakuba* and in *D. simulans* (Figure 5A). However, few genes ($\sim 11\%$) are duplicated in both species and only a handful have been identified in *D. simulans*, *D. yakuba*, and *D. melanogaster* (Figure 5B). Moreover, none of the high frequency variants in the in *D. yakuba* and *D. simulans* capture orthologous sequences. Hence, in spite of parallel selective pressures on rapidly evolving phenotypes, there is little parallel recruitment of the same genetic solutions with respect to duplication. Given the limited genomic span of standing variation in the population (Table S7), and low rates of new mutation (Figure 6, Table

1), as well as the low frequency of a large fraction of variants, parallel fixation of tandem duplications in the same genes will be extremely rare even among genera with large effective population sizes facing similar selective pressures.

Convergence depends on the wait time of new mutations to enter populations and establish selective sweeps. We show that the average wait time for a new mutation given a selection coefficient of $s = 0.01$ is hundreds of years for SNPs. Here, we find that tandem duplications display signals of reduced heterozygosity in the surrounding regions as well as an association with gene ontologies indicative of rapidly evolving phenotypes, and an overrepresentation of shared tandem duplicates across species for specific genes given the limits of standing variation, consistent with widespread adaptation through tandem duplication. However, the average wait time for a deterministic sweep to establish in a population will be thousands of years for tandem duplications and tens of thousands of years for chimeric genes given a modest selection coefficient of $s = 0.01$. Such strongly selected sites are expected to be rare throughout the genome and beneficial mutations are likely to appear less often than the actual mutation rate (Jensen, Thornton and Andolfatto 2008, Andolfatto 2007). Thus, these wait times given strong selection provide a lower bound to the wait time for a selected sweep. We therefore expect that mutation will severely limit evolution through whole gene duplication and chimera formation, to the extent that adaptation depends on tandem duplications, the ability of organisms to adapt to changing environments will be hindered by a lack of variation. Thus, even when a given tandem duplication is needed for adaptation, we expect that the limits of mutation will lead to low

levels of convergence and scarcity of shared genetic solutions.

Duplicate genes and rapidly evolving phenotypes

Both *D. simulans* and *D. yakuba* have an overabundance of genes with ontology classifications involved in immune function, chemosensory processing or response, and drug and toxin metabolism (Table S5). Furthermore the instance of independent duplications confirm a bias toward chemosensory receptors, chorion development and oogenesis, as well as immune response (Rogers et al. 2014). These phenotypes are strongly associated with rapid evolution due to host-parasite interactions, predator-prey coevolution, and sexual conflict (Lazarro and Clark 2012, Beckerman et al. 2013, Ffrench-Constant, Daborn and Le Goff 2004, Panhuis, Clark and Swanson 2006). Previous work has observed similar bias toward rapid amino acid substitutions in olfactory genes, and chitin cuticle genes in *D. melanogaster* and *D. simulans* (Begun et al. 2007), selection for gene family evolution in and selection for toxin resistance is common in *D. melanogaster* (Schmidt et al. 2010, Ffrench-Constant, Daborn and Le Goff 2004) suggesting that associated phenotypes may be under widespread selection in multiple species.

Host pathogen systems as well as arms races in pesticide and toxin resistance operate under Red Queen dynamics in which conflicts between organisms result in repeated selective sweeps (Van Valen 1973). Organisms that lack the genetic means to adapt to rapidly changing systems will be at a distinct disadvantage in the face of selective events. Additionally, the overrepresentation of duplications in cytochromes and drug or

toxin metabolism genes confirms rapid evolution in copy number seen in comparison of reference genomes (Drosophila Twelve Genomes Consortium 2007) as well as recent studies of insecticide resistance and viral resistance in natural populations (Schmidt et al. 2010, Bass and Field 2011, Magwire et al. 2011). Large amounts of divergence driven by selection among non-synonymous sites and UTRs in *D. simulans* (Haddrill, Bachtrog and Andolfatto 2008) and high rates of adaptive substitutions (Andolfatto, Wong and Bachtrog 2011b, Begun et al. 2007) point to widespread selective pressures acting in *D. simulans*, and it is likely that these same pressures influence the current diversity and frequency of copy number variants. If rapidly evolving systems rely heavily on complex mutations or if selection coefficients are modest, profiles of standing variation will place strong limits on outcomes in response to selection.

Shifting selective pressures such as those found in rapidly evolving systems or gross ecological change require a pool of genetic variation to facilitate adaptation. We observe standing variation and mutational profiles that will limit evolutionary trajectories and would expect these limits to be even more severe for rapidly evolving phenotypes. Repeated sweeps are expected to purge genetic and phenotypic diversity, and recovering such diversity after sweeps can take thousands of generations (Kaplan, Hudson and Langley 1989). Thus, during rapid evolution, selection will potentially purge diversity that is needed for subsequent steps in the adaptive walk. Hence, although duplications are key players in rapid evolution, their limited rates of formation combined with low frequencies due to commonly detrimental impacts will hinder evolutionary outcomes or force alternative adaptive trajectories precisely

when variation is urgently needed. Moreover, large numbers of duplicates are low-frequency, suggesting that detrimental impacts further limit standing variation. Thus, we conclude that the available substrate of tandem duplications and profiles of standing variation will define evolutionary outcomes in *Drosophila* and other multicellular eukaryotes.

Materials and Methods

Tandem duplications

Tandem duplications were identified using paired-end Illumina sequencing of genomic DNA for 20 strains of *D. yakuba* and 20 strains of *D. simulans* as well as the reference genome of each species as described in Rogers et al. 2014. The dataset describes derived, segregating tandem duplications that span 25 kb or less. These sequences exclude ancestral duplications as well as putative duplications in the resequenced reference genomes. The resulting list of variants describes segregating variation for newly formed tandem duplicates across the full genome in these two species of non-model *Drosophila*.

Identifying duplicated coding sequence

Tandem duplications were previously identified using a combination of paired end read mapping and coverage changes in 20 isofemale lines of *D. yakuba* and 20 isofemale lines in *D. simulans* generated via 9-12 generations of sibling mating from wild-caught flies. We sequenced 10 isofemale lines of *D. yakuba* from Nairobi, Kenya, and 10 isofemale lines from

Nguti, Cameroon as well as 10 isofemale lines of *D. simulans* from Nairobi, Kenya and 10 isofemale lines from Madagascar. Duplications were identified through divergently oriented reads and coverage changes in comparison to reference genomes. We identify 1415 tandem duplications in *D. simulans* and 975 tandem duplications segregating in *D. yakuba* that span 845 different gene sequences in *D. yakuba* and 478 different gene sequences in *D. simulans* (Rogers et al. 2014). Gene duplications were defined as any divergent read calls whose maximum span across all lines overlaps with the annotated CDS coordinates. *D. yakuba* CDS annotations were based on flybase release *D. yakuba* r.1.3. Gene annotations for the recent reassembly of the *D. simulans* reference were produced by aligning all *D. melanogaster* CDS sequences to the *D. simulans* reference in a tblastx. Percent coverage of the CDS was defined based on the portion of the corresponding genomic sequence from start to stop that was covered by the maximum span of divergent read calls across all strains. Using the representation of gene sequences in *D. yakuba* of $\frac{845}{16082}$ we use a binomial test to calculate the likelihood of 56 shared variants among the 478 genes duplicated in *D. simulans*.

Frequency of unsampled alleles

We also estimated the frequency distribution for alleles that are absent among our 20 strains, according to a Bayesian binomial model. Assuming that sampling follows a binomial model that is dependent upon the allele frequency p , the probability that a variant is present at frequency p given that it is not observed is as follows:

$$P(p|absent) = \frac{p(absent|p)*p(p)}{\int_0^1 p(absent|p)*p(p)dp}.$$

Assuming a uniform distribution on p :

$$P(p|absent) = \frac{p(absent|p)}{\int_0^1 p(absent|p)dp.}$$

$$P(p|absent) = \frac{(1-p)^{20}}{\int_0^1 (1-p)^{20}dp.}$$

$$P(p|absent) = 21(1-p)^{20}$$

95% lower CI defined as:

$$\int_0^x 21(1-p)^{20}dp = 0.95$$

$$1 - (1-x)^{21} = 0.95$$

And therefore the 95% one-sided lower CI is $x \leq 0.132946$ whereas the 50% one-sided lower CI will be $x \leq 0.0325$. Placing a uniform prior on p will bias estimates toward higher frequency variants, thereby placing a conservative upper bound on allele frequencies.

Estimated number of segregating tandem duplications

We compared the estimated total number of duplications expected in a population to estimates of diversity based on our sample of 20 strains, correcting S for a 3.9% false positive rate (Table S6). Under a standard coalescent model (Wakeley 2009, Ewens 1974, Watterson 1975):

$$E[S_{population}] = \frac{S_{sample}}{a_{sample}} * a_{population}$$

Where a in a sample of size n (in this case $n=20$):

$$a_{sample} = \sum_{i=0}^{n-1} \frac{1}{i}$$

$$a_{population} = \sum_{i=0}^{2N_e} \frac{1}{i}$$

When $2N_e$ is large:

$$\sum_{i=0}^{2N_e} \frac{1}{i} \approx \theta(\ln(2N_e) + 0.57722)$$

Hence:

$$E[S_{population}] = \frac{S_{sample}}{a_{20}} * (\ln(2N_e) + 0.57722)$$

We can use similar methods to estimate the variance in the number of segregating sites in the population.

$$Var[S_{population}] = \theta \sum_{i=0}^{2N_e} \frac{1}{i} + \theta^2 \sum_{i=0}^{2N_e} \frac{1}{i^2}$$

When $2N_e$ is large:

$$\sum_{i=0}^{2N_e} \frac{1}{i^2} \approx \frac{\pi^2}{6}.$$

$$Var[S_{population}] = \theta(\ln(2N_e) + 0.57722) + \theta^2 \frac{\pi^2}{6}$$

Gene Ontology

Overrepresented functional categories were identified using DAVID gene ontology software with an EASE threshold of 1.0. as previously described (Rogers et al. 2014). We observe several functional categories indicative of rapid evolution that are shared between the two species (Table S5). As a comparison, we selected a random subset of 845 genes for *D. yakuba* and 478 genes from *D. simulans*, and performed ontology analysis for a comparison. In *D. yakuba*, male courtship, GTPase enzymes, and alt splicing had significant group EASE thresholds whereas *D. simulans* showed marginal values on esterases, tracheal development, neurodevelopment, lipid metabolism, hormone receptors, cell communication and growth and starvation. Nothing has an EASE of 1.5 in either species similar to the values observed in Table S5. There is no agreement in functional categories for the two species in the random

subsets. Thus, we would suggest that the convergence across species and the associations with rapid evolution are not the product of sampling errors.

Proportion of the genome represented by segregating duplicates

To determine the number of duplications necessary to span the full range of the genome, we simulated chromosomes with a length determined by the number of base pairs with non-zero coverage in our reference strain. We then simulated random draws from the distribution of duplication lengths for each chromosome, placing duplication start sites at random and recorded the number of duplications necessary to cover 10%, 25%, 50%, and 90% of sequence length for each chromosome in each trial. Simulations were repeated for 1000 trials for each chromosome.

These simulations do not account for mutational biases that might result in clustering of duplications in particular regions while other regions remain static, nor do they require that new duplications reach an appreciable frequency so that they are immune to stochastic loss through genetic drift. They do not require that duplications capture sufficient sequence to have functional impacts or require that breakpoints not disrupt known functional elements. Furthermore, simulating individual chromosomes separately decreases the likelihood of resampling particular sites thereby lowering the estimated number of duplications needed to cover the entire genome. Hence, these estimates put a highly conservative lower bound on the minimum number of mutations necessary to capture the full genomic sequence.

To estimate the expected proportion of the genome spanned by all duplicates in the

population, we resampled 6700 duplicates from the observed size distribution of *D. yakuba* with replacement and 4000 duplicates from the observed size distribution of *D. simulans*, placing duplications at random positions across the chromosome. We performed 100 replicates of sampling and report the mean across all replicates for each species. In *D. simulans* we observe one case with 19 independent whole gene duplications of a single ORF (Rogers et al. 2014), suggesting up to 1000-fold variation in mutation rates over the genome average. Estimates of population level variation and genome wide mutation rates ignore mutation rate variation where some regions may be highly prone to duplications whereas others remain static, which would reduce likelihood of unobserved tandem duplications outside of mutational hotspots. Hence, these estimates represent a lower bound on the number of duplications necessary to span the entire genome.

Mutation rates and wait times for duplicates

We estimate average heterozygosity (θ_π) and mutation rates (μ) per gene for *D. yakuba* and *D. simulans* for gene duplications that capture at least 90% of gene sequence (in agreement with previous estimates (Rogers, Bedford and Hartl 2009)), for genes that recruit non-coding sequence, and for chimeric genes. Heterozygosity estimates used to calculate mutation rates were corrected for ascertainment bias (see SI Text) and excluded genes that were captured by 4 or more independent mutations, a signal of hotspots and mutation rate heterogeneity. Heterozygosity per gene is estimated given 16,082 gene sequences in *D. yakuba* and 10,786 coding sequences in *D. simulans* (Table 1). Given estimates of θ_π , we estimate the probability

of adaptation from standing variation under an additive genetic model for neutral variants, $P_{sgv} = e^{-\theta_{\pi} * \ln(1+2N_e s)}$ (Hermisson and Pennings 2005) and the time to establishment (T_e) of a deterministic sweep from new mutations, such that new mutants escape the stochastic forces of drift, $T_e = \frac{1}{\theta_{\pi} s}$ (Maynard Smith 1971, Gillespie 1991). These estimates are provided for two strong selection coefficients of $s = 0.01$ typical of what is observed in *Drosophila* and $s = 0.20$ modeling abnormally strong selection on a single locus consistent with Karasov, Messer and Petrov (2010). Estimates assume that a given site of interest is under strong selection and that the beneficial mutation rate is equal to the mutation rate per site per generation providing an upper limit on the ability of new mutations to facilitate adaptation. In reality, strongly selected sites will be rare throughout the genome (Jensen, Thornton and Andolfatto 2008, Andolfatto 2007) and these wait times given strong selection will not reflect the expected number of selective sweeps throughout the genome.

Additional methods

Further description of methods including description of intronic SNPs, analysis of population structure, residual heterozygosity, calculation of N_e , and correction for ascertainment bias are available in SI Text. All data files are available via <http://molpopgen.org/Data> and <http://www.github.com/ThorntonLab/DrosophilaPopGenData-Rogers2014>. Aligned bam files were deposited in the National Institutes of Health Short Read Archive under accession numbers SRP040290 and SRP029453. Sequenced stocks were deposited in the University of California, San Diego (UCSD) stock center with stock numbers 14021-0261.38- 14021-0261.51

and 14021-0251.293 - 14021-0251.311.

Acknowledgements

The authors would like to thank Nigel F. Delaney, Elizabeth G. King, Anthony D. Long, and Alexis S. Harrison, and Trevor Bedford for helpful discussions as well as three anonymous reviewers whose comments substantially improved the manuscript. RLR is supported by NIH Ruth Kirschstein National Research Service Award F32-GM099377. Research funds were provided by NIH grant R01-GM085183 to KRT and R01-GM083228 to PA. All sequencing was performed at the UC Irvine High Throughput Genomics facility, which is supported by the National Cancer Institute of the National Institutes of Health under Award Number P30CA062203. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. RLR, JMC, KRT, and PA performed analyses. LS generated Illumina sequencing libraries. TTH provided gene annotations for *D. simulans*. RLR, JMC, PA and KRT designed experiments and analyses.

Bibliography

- Andolfatto P. 2007. Hitchhiking effects of recurrent beneficial amino acid substitutions in the *Drosophila melanogaster* genome. *Genome Res.* 17:1755–1762.
- Andolfatto P, Wong KM, Bachtrog D. 2011a. Effective population size and the efficacy of selection on the X chromosomes of two closely related *Drosophila* species. *Genome Biol Evol.* 3:114–128.
- Andolfatto P, Wong KM, Bachtrog D. 2011b. Effective population size and the efficacy of selection on the X chromosomes of two closely related *Drosophila* species. *Genome Biol Evol.* 3:114–128.
- Bachtrog D, Thornton K, Clark A, Andolfatto P. 2006. Extensive introgression of mitochondrial DNA relative to nuclear genes in the *Drosophila yakuba* species group. *Evolution.* 60:292–302.
- Bass C, Field LM. 2011. Gene amplification and insecticide resistance. *Pest Manag. Sci.* 67:886–890.
- Beckerman AP, de Roij J, Dennis SR, Little TJ. 2013. A shared mechanism of defense against predators and parasites: chitin regulation and its implications for life-history theory. *Ecol Evol.* 3:5119–5126.
- Begun DJ, Holloway AK, Stevens K, et al. (13 co-authors). 2007. Population genomics: whole-genome analysis of polymorphism and divergence in *Drosophila simulans*. *PLoS Biol.* 5:e310.
- Brodie ED. 2010. Convergent evolution: pick your poison carefully. *Curr. Biol.* 20:R152–154.
- Burke MK, Dunham JP, Shahrestani P, Thornton KR, Rose MR, Long AD. 2010. Genome-wide analysis of a long-term evolution experiment with *Drosophila*. *Nature.* 467:587–590.
- Cardoso-Moreira M, Emerson JJ, Clark AG, Long M. 2011. *Drosophila* duplication hotspots are associated with late-replicating regions of the genome. *PLoS Genet.* 7:e1002340.
- Chan YF, Marks ME, Jones FC, et al. (16 co-authors). 2010. Adaptive evolution of pelvic reduction in sticklebacks by recurrent deletion of a *Pitx1* enhancer. *Science.* 327:302–305.

- Charlesworth B, Morgan MT, Charlesworth D. 1993. The effect of deleterious mutations on neutral molecular variation. *Genetics*. 134:1289–1303.
- Chen L, DeVries AL, Cheng CH. 1997. Convergent evolution of antifreeze glycoproteins in Antarctic notothenioid fish and Arctic cod. *Proc. Natl. Acad. Sci. U.S.A.* 94:3817–3822.
- Colosimo PF, Hosemann KE, Balabhadra S, Villarreal G, Dickson M, Grimwood J, Schmutz J, Myers RM, Schluter D, Kingsley DM. 2005. Widespread parallel evolution in sticklebacks by repeated fixation of Ectodysplasin alleles. *Science*. 307:1928–1933.
- Conant GC, Wolfe KH. 2008. Turning a hobby into a job: how duplicated genes find new functions. *Nat. Rev. Genet.* 9:938–950.
- Coop G, Ralph P. 2012. Patterns of neutral diversity under general models of selective sweeps. *Genetics*. 192:205–224.
- Cridland JM, Thornton KR. 2010. Validation of rearrangement break points identified by paired-end sequencing in natural populations of *Drosophila melanogaster*. *Genome Biol. Evol.* 2:83–101.
- Dopman EB, Hartl DL. 2007. A portrait of copy-number polymorphism in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. U.S.A.* 104:19920–19925.
- Drosophila Twelve Genomes Consortium. 2007. Evolution of genes and genomes on the *Drosophila* phylogeny. *Nature*. 450:203–218.
- Emerson JJ, Cardoso-Moreira M, Borevitz JO, Long M. 2008. Natural selection shapes genome-wide patterns of copy-number polymorphism in *Drosophila melanogaster*. *Science*. 320:1629–1631.
- Ewens WJ. 1974. A note on the sampling theory for infinite alleles and infinite sites models. *Theor. Popul. Biol.* 6:143–148.
- Eyre-Walker A, Keightley PD, Smith NG, Gaffney D. 2002. Quantifying the slightly deleterious mutation model of molecular evolution. *Mol. Biol. Evol.* 19:2142–2149.
- French-Constant RH, Daborn PJ, Le Goff G. 2004. The genetics and genomics of insecticide resistance. *Trends Genet.* 20:163–170.
- Gillespie JH. 1991. *The Causes of Molecular Evolution*, Oxford University Press, chapter 5, p. 232.
- Gould SJ, Lewontin RC. 1979. The spandrels of San Marco and the Panglossian paradigm: a critique of the adaptationist programme. *Proc. R. Soc. Lond., B, Biol. Sci.* 205:581–598.
- Haddrill PR, Bachtrog D, Andolfatto P. 2008. Positive and negative selection on noncoding DNA in *Drosophila simulans*. *Mol. Biol. Evol.* 25:1825–1834.

- Haldane JBS. 1927. A mathematical theory of natural and artificial selection V: Selection and mutation. *Proceedings of the Cambridge Philosophical Society*. 23:838–844.
- Hermisson J, Pennings PS. 2005. Soft sweeps: molecular population genetics of adaptation from standing genetic variation. *Genetics*. 169:2335–2352.
- Hu TT, Eisen MB, Thornton KR, Andolfatto P. 2012. A second generation assembly of the *Drosophila simulans* genome provides new insights into patterns of lineage-specific divergence. *Genome Res.* .
- Jensen JD, Thornton KR, Andolfatto P. 2008. An approximate bayesian estimator suggests strong, recurrent selective sweeps in *Drosophila*. *PLoS Genet.* 4:e1000198.
- Kaplan NL, Hudson RR, Langley CH. 1989. The “hitchhiking effect” revisited. *Genetics*. 123:887–899.
- Karasov T, Messer PW, Petrov DA. 2010. Evidence that adaptation in *Drosophila* is not limited by mutation at single sites. *PLoS Genet.* 6:e1000924.
- Katju V, Lynch M. 2006. On the formation of novel genes by duplication in the *Caenorhabditis elegans* genome. *Molecular Biology and Evolution*. 23:1056–1067.
- Khadjeh S, Turetzek N, Pechmann M, Schwager EE, Wimmer EA, Damen WG, Prpic NM. 2012. Divergent role of the Hox gene *Antennapedia* in spiders is responsible for the convergent evolution of abdominal limb repression. *Proc. Natl. Acad. Sci. U.S.A.* 109:4921–4926.
- Kreitman M. 1983. Nucleotide polymorphism at the alcohol dehydrogenase locus of *Drosophila melanogaster*. *Nature*. 304:412–417.
- Lazarro B, Clark A. 2012. *Rapidly Evolving Genes and Genetic Systems*, Oxford: Oxford University Press, chapter Rapid evolution of innate immune response genes.
- Lemeunier F, Ashburner MA. 1976. Relationships within the melanogaster species subgroup of the genus *Drosophila* (Sophophora). II. Phylogenetic relationships between six species based upon polytene chromosome banding sequences. *Proc. R. Soc. Lond., B, Biol. Sci.* 193:275–294.
- Llopart A, Lachaise D, Coyne JA. 2005. Multilocus analysis of introgression between two sympatric sister species of *Drosophila*: *Drosophila yakuba* and *D. santomea*. *Genetics*. 171:197–210.
- Lynch M. 2007. *The Origins of Genome Architecture*, Sunderland, Mass.: Sinauer Associates, p. 494.
- Lynch M, Conery JS. 2000. The evolutionary fate and consequences of duplicate genes. *Science*. 290:1151–1155.

- Lynch M, Conery JS. 2003. The evolutionary demography of duplicate genes. *Journal of Structural and Functional Genomics*. 3:35–44.
- Magwire MM, Bayer F, Webster CL, Cao C, Jiggins FM. 2011. Successive increases in the resistance of *Drosophila* to viral infection through a transposon insertion followed by a Duplication. *PLoS Genet*. 7:e1002337.
- Martin A, Papa R, Nadeau NJ, et al. (11 co-authors). 2012. Diversification of complex butterfly wing patterns by repeated regulatory evolution of a Wnt ligand. *Proc. Natl. Acad. Sci. U.S.A.* 109:12632–12637.
- Maynard Smith J. 1971. What use is sex? *Journal of theoretical biology*. 30:319–335.
- Moxon E, Lenski R, Rainey P. 1998. Adaptive evolution of highly mutable loci in pathogenic bacteria. *Perspectives in Biology and Medicine*. 42:154–155.
- Nielsen R, Bustamante C, Clark AG, et al. (13 co-authors). 2005. A scan for positively selected genes in the genomes of humans and chimpanzees. *PLoS Biol*. 3:e170.
- Ohno S. 1970. *Evolution by gene duplication*. London: George Alien & Unwin Ltd. Berlin, Heidelberg and New York: Springer-Verlag.
- Orozco-terWengel P, Kapun M, Nolte V, Kofler R, Flatt T, Schloetterer C. 2012. Adaptation of *drosophila* to a novel laboratory environment reveals temporally heterogeneous trajectories of selected alleles. *Molecular ecology*. 21:4931–4941.
- Panhuis TM, Clark NL, Swanson WJ. 2006. Rapid evolution of reproductive proteins in abalone and *Drosophila*. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 361:261–268.
- Presgraves DC. 2008. Sex chromosomes and speciation in *Drosophila*. *Trends Genet*. 24:336–343.
- Riehle MM, Bennett AF, Long AD. 2001. Genetic architecture of thermal adaptation in *Escherichia coli*. *Proc. Natl. Acad. Sci. U.S.A.* 98:525–530.
- Rogers RL, Bedford T, Hartl DL. 2009. Formation and longevity of chimeric and duplicate genes in *Drosophila melanogaster*. *Genetics*. 181:313–322.
- Rogers RL, Cridland JM, Shao L, Hu TT, Andolfatto P, Thornton KR. 2014. Landscape of Standing Variation for Tandem Duplications in *Drosophila yakuba* and *Drosophila simulans*. *Mol. Biol. Evol.* 31:1750–1766.
- Rogers RL, Hartl DL. 2012. Chimeric genes as a source of rapid evolution in *Drosophila melanogaster*. *Mol. Biol. Evol.* 29:517–529.
- Schmidt JM, Good RT, Appleton B, et al. (11 co-authors). 2010. Copy number variation and transposable elements feature in recent, ongoing adaptation at the *Cyp6g1* locus. *PLoS Genet*. 6:e1000998.

- Schrider DR, Stevens K, Cardeno CM, Langley CH, Hahn MW. 2011. Genome-wide analysis of retrogene polymorphisms in *Drosophila melanogaster*. *Genome Res.* 21:2087–2095.
- Shapiro MD, Summers BR, Balabhadra S, Aldenhoven JT, Miller AL, Cunningham CB, Bell MA, Kingsley DM. 2009. The genetic architecture of skeletal convergence and sex determination in ninespine sticklebacks. *Curr. Biol.* 19:1140–1145.
- Smith JM, Haigh J. 1974. The hitch-hiking effect of a favourable gene. *Genet. Res.* 89:391–403.
- Stern DL. 2013. The genetic causes of convergent evolution. *Nat. Rev. Genet.* 14:751–764.
- Tajima F. 1983. Evolutionary relationship of DNA sequences in finite populations. *Genetics.* 105:437–460.
- Tajima F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics.* 123:585–595.
- Tamura K, Subramanian S, Kumar S. 2004. Temporal patterns of fruit fly (*Drosophila*) evolution revealed by mutation clocks. *Mol. Biol. Evol.* 21:36–44.
- Tanaka K, Barmina O, Kopp A. 2009. Distinct developmental mechanisms underlie the evolutionary diversification of *Drosophila* sex combs. *Proc. Natl. Acad. Sci. U.S.A.* 106:4764–4769.
- Tenaillon O, Rodriguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD, Gaut BS. 2012. The molecular diversity of adaptive convergence. *Science.* 335:457–461.
- Thornton K, Long M. 2002. Rapid divergence of gene duplicates on the *Drosophila melanogaster* X chromosome. *Mol. Biol. Evol.* 19:918–925.
- Thornton K, Long M. 2005. Excess of amino acid substitutions relative to polymorphism between X-linked duplications in *Drosophila melanogaster*. *Mol. Biol. Evol.* 22:273–284.
- Van Valen L. 1973. A new evolutionary law. *Evolutionary theory.* 1:1–30.
- Wakeley J. 2009. *Coalescent Theory: An Introduction*, Roberts & Company Publishers, p. 97.
- Watterson GA. 1975. On the number of segregating sites in genetical models without recombination. *Theor Popul Biol.* 7:256–276.
- Wittkopp PJ, Williams BL, Selegue JE, Carroll SB. 2003. *Drosophila* pigmentation evolution: divergent genotypes underlying convergent phenotypes. *Proc. Natl. Acad. Sci. U.S.A.* 100:1808–1813.

- Woods R, Schneider D, Winkworth CL, Riley MA, Lenski RE. 2006. Tests of parallel molecular evolution in a long-term experiment with *Escherichia coli*. *Proceedings of the National Academy of Sciences*. 103:9107–9112.
- Zhen Y, Aardema ML, Medina EM, Schumer M, Andolfatto P. 2012. Parallel molecular evolution in an herbivore community. *Science*. 337:1634–1637.
- Zhou Q, Zhang G, Zhang Y, Xu S, Zhao R, Zhan Z, Li X, Ding Y, Yang S, Wang W. 2008. On the origin of new genes in *Drosophila*. *Genome Research*. 18:1446–1455.
- Zichner T, Garfield DA, Rausch T, Stutz AM, Cannavo E, Braun M, Furlong EE, Korbel JO. 2013. Impact of genomic structural variation in *Drosophila melanogaster* based on population-scale sequencing. *Genome Res*. 23:568–579.

Table 1: Establishment of Sweeps in *D. yakuba* and *D. simulans*.

Species		Intron SNPs	Whole Gene	Recruit	Chimera
<i>D. yakuba</i>	μ	5.8×10^{-9}	1.17×10^{-9}	3.46×10^{-10}	3.70×10^{-10}
	θ_π	0.0138	0.00277	0.00082	0.00088
	P_{sgv} , s=0.01	12.1%	2.23%	0.67%	0.71%
	P_{sgv} , s=0.20	15.7%	3.05%	0.91%	0.97%
	T_e , s=0.01	7270	36,000	122,000	114,000
	T_e , s=0.20	364	1,800	6,087	5,704
Species		Intron SNPs	Whole Gene	Recruit	Chimera
<i>D. simulans</i>	μ	5.8×10^{-9}	6.03×10^{-10}	2.42×10^{-10}	8.52×10^{-11}
	θ_π	0.0280	0.00291	0.00117	0.00041
	P_{sgv} , s=0.01	24.6%	2.56%	1.04%	0.37%
	P_{sgv} , s=0.20	30.1%	3.41%	1.38%	0.49%
	T_e , s=0.01	3580	34,400	85,700	243,000
	T_e , s=0.20	179	1,720	4,290	12,100

P_{sgv} from Hermisson and Pennings (2005) estimates the likelihood of adaptation from standing genetic variation under an additive model assuming neutral variation.

T_e (Gillespie 1991 and Maynard-Smith 1971) estimates the average time until establishment of a selective sweep from a new mutation in generations given that a site is under strong selection with beneficial mutation rate equal to θ_π . Estimates provide a lower bound on T_e .




Figure 1: SFS for tandem duplications in *D. yakuba* and *D. simulans*, corrected for ascertainment bias. A. Site frequency spectra on the autosomes (black) and on the X (grey) in *D. yakuba*. B. SFS on the autosomes (black) and on the X (grey) in *D. simulans*. C. SFS for X-linked intronic SNPs (black) and duplicates (grey). The excess of high frequency variants on the X in *D. simulans* suggests widespread selection for tandem duplicates on the *D. simulans* X.




Figure 2: Diversity (θ_π) as a function of distance from new mutations in *D. yakuba* for putatively neutral intronic SNPs (black) and for tandem duplications (red) by chromosome with lowess smoothing. Duplicates show a reduction in diversity approaching duplications, whereas neutral SNPs show no reduction in diversity. Plots exclude centromeric regions and the 4th chromosome which have atypical nucleotide diversity.




Figure 3: Nucleotide diversity, θ_π as a function of distance from new mutations in *D. simulans* for putatively neutral intronic SNPs (black) and for tandem duplications (red) by chromosome with lowest smoothing. Duplicates show a reduction in mean diversity approaching duplications, whereas neutral SNPs show no reduction in diversity. Plots exclude centromeric regions and the 4th chromosome which have atypical nucleotide diversity. Chromosome 3L is strongly affected by a cluster of duplications at roughly 8.5Mb, which is excluded from the plot.




Figure 4: Histogram of nucleotide diversity, θ_π , (A) For Intergenic mutations (yellow) and duplications that capture gene sequences but do not create chimeric constructs (blue) in *D. yakuba*. (B) For Intergenic mutations (yellow) and duplications that create chimeric genes (red) in *D. yakuba*. (C) For Intergenic mutations (yellow) and duplications that capture gene sequences but do not create chimeric constructs (blue) in *D. simulans*. (D) For Intergenic mutations (yellow) and duplications that create chimeric genes (red) in *D. simulans*.




Figure 5: A) Gene ontology classes overrepresented by species among singly duplicated genes or among multiply duplicated genes. B) Number of genes duplicated by species. Most variants are species specific, with small numbers of parallel duplication of orthologs across species.




Figure 6: Genomewide population mutation rates for all duplications (θ), population sizes (N_e), and per gene mutation rates (μ) for gene structures produced by duplication by species. Low mutation rates and mutation limited evolution leads to low levels of parallel recruitment of tandem duplications.

Supporting Information

Identifying duplicated coding sequence

Tandem duplications were previously identified using a combination of paired end read mapping and coverage changes in 20 isofemale lines of *D. yakuba* and 20 isofemale lines in *D. simulans* generated via 9-12 generations of sibling mating from wild-caught flies. We sequenced 10 isofemale lines of *D. yakuba* from Nairobi, Kenya, and 10 isofemale lines from Nguti, Cameroon as well as 10 isofemale lines of *D. simulans* from Nairobi, Kenya and 10 isofemale lines from Madagascar. Duplications were identified through divergently oriented reads and coverage changes in comparison to reference genomes. We identify 1415 tandem duplications in *D. simulans* and 975 tandem duplications segregating in *D. yakuba* that span 845 different gene sequences in *D. yakuba* and 478 different gene sequences in *D. simulans* (Rogers et al. 2014). Gene duplications were defined as any divergent read calls whose maximum span across all lines overlaps with the annotated CDS coordinates. *D. yakuba* CDS annotations were based on flybase release *D. yakuba* r.1.3. Gene annotations for the recent reassembly of the *D. simulans* reference were produced by aligning all *D. melanogaster* CDS sequences to the *D. simulans* reference in a tblastx. Percent coverage of the CDS was defined based on the portion of the corresponding genomic sequence from start to stop that

was covered by the maximum span of divergent read calls across all strains. Using the representation of gene sequences in *D. yakuba* of $\frac{845}{16082}$ we use a binomial test to calculate the likelihood of 56 shared variants among the 478 genes duplicated in *D. simulans*.

Intronic SNPs

In order to produce a neutral proxy for sequence change in each species, we identified SNPs for short introns 100 bp or less, focusing on sites 8-30 which are generally subject to little constraints (Halligan and Keightley 2006, Parsch et al. 2010, Clemente and Vogl 2012). Reads containing indels were re-aligned using GATK (McKenna et al. 2010). SNPs were identified across strains using samtools v1.18 mpileup (Li et al. 2009) disabling probabilistic realignment (-B) and outputting genotype likelihoods in BCF format (-g). The resulting BCF used to create a VCF using bcftools, calling bases using Bayesian inference (-c) calling genotypes per sample (-g) with a scaled mutation rate of 1% (-t .01) under a haploid model (ploidy=1). SNPs were required to have minimum Illumina coverage depth of 20 reads, maximum coverage of 250 reads, $MQ \geq 20$, and $GQ \geq 30$ and invar $GQ \geq 40$. We excluded SNPs identified in the reference, which are indicative of either assembly errors or residual heterozygosity. We performed hierarchical cluster analysis in R using all SNPs by chromosome to evaluate population structure.

The ancestral state for each SNP was established through comparison with the nearest sequenced reference genome as an outgroup, *D. erecta* for *D. yakuba* sequences and *D. melanogaster* for *D. simulans* sequences. Orthologs between each species and its outgroup

were identified using reciprocal best hit criteria in a BLASTn at an E-value cutoff of 10^{-5} . Full gene sequences for each ortholog were then aligned using clustalw, keeping only genes which aligned with 85% or greater nucleotide identity. Divergence between the two species, $Div_{x,y}$, was defined based on alignments of intronic sites from bases 8-30 between each species and the outgroup reference genome, excluding gapped sequences, for aligned orthologs with 85% or nucleotide identity. The ancestral state was defined based on the corresponding sequence in the outgroup genome. We excluded sites where the outgroup reference was in disagreement with both the *D. yakuba* reference and *D. yakuba* SNPs, as well as triallelic SNPs, sites with reference sequence of ‘N’, or SNPs identified in the VCF for the reference, suggesting inaccuracies in reference assembly or residual heterozygosity in the reference. These resulted in a total of 7158 intronic SNPs in *D. yakuba* and 5504 intronic SNPs in *D. simulans*. The resulting unfolded SFS was then corrected for the probability of independent mutations in both reference genomes leading to incorrect inference of the ancestral state.

Given net divergence $D_{net} = Div_{x,y} - \pi_x$, the probability of identical independent mutations occurring in the outgroup reference genome is reflected by either the probability of an independent transition (ts) at the site of a transition mutation, or by 1/2 the probability of a transversion (tv) at the site of a transversion polymorphism. Thus,

$$k = [(\frac{\kappa}{2 + \kappa})^2 + \frac{1}{2}(\frac{2}{2 + \kappa})^2]D_{net} \quad (1)$$

Empirically, in *Drosophila* $\kappa = \frac{ts}{tv} = 2$. Thus, $k = \frac{3}{8}D_{net}$.

The unfolded SFS for intronic sites was corrected for the likelihood of independent

mutations in the reference, k . The probability of independent mutations occurring in both genomes is equal to the probability of either two independent transitions or two independent transversions occurring in both genomes. We calculated π_x as the average heterozygosity per intronic site.

Given a likelihood of independent identical mutations of $k = \frac{3}{8}D_{net}$ (See Text S1).

$$S_{i,obs} = E[S_i] - E[S_i](k) + E[S_{n-i}](k) \quad (2)$$

$$S_{n-i,obs} = E[S_{n-i}] - E[S_{n-i}](k) + E[S - i](k) \quad (3)$$

Substituting Equation 3 into Equation 2, we obtain

$$E[S_i] = \frac{S_{i,obs}(1 - k) - S_{n-i,obs}(k)}{1 - 2k} \quad (4)$$

Correcting Duplicates for Ascertainment Bias

Tandem duplications, unlike SNPs, cannot be identified using paired-end reads in individual strains except through comparison to the reference genome. Moreover, variants that are segregating at high frequency in populations are substantially more likely to be present in the reference, and therefore are substantially less likely to be identified in sample strains (Emerson et al. 2008). We corrected site frequency spectra according to the model developed by Emerson et al. (Emerson et al. 2008).

$$x_i = \frac{y_{i \frac{n}{n-i}}}{\sum_{i=1}^{n-2} y_{i \frac{n}{n-i}}} \quad (5)$$

Here, x_i is the true proportion of alleles at frequency i in the population, and y_i is the observed proportion of alleles at frequency i in a sample of n strains (here 21). The correction for ascertainment bias lowers estimates of the proportion found at low frequencies and increases estimates of the proportion at high frequency. For estimates of population site frequency spectra, we removed all variants with divergently oriented reads in the reference strain, as these would not be identified in an accurately annotated reference.

Residual heterozygosity

Some isofemale lines contained regions of residual heterozygosity in spite of over 10 generations of inbreeding in the lab. To detect regions of residual heterozygosity, we called SNPs as above under a diploid model. Segments with residual heterozygosity were detected using an HMM (HMM; <http://cran.r-project.org/web/packages/HMM/>).

Prior probabilities on states were set as:

$$\pi = [0.5 \quad 0.5]$$

Transition probabilities were set to:

$$T = \begin{bmatrix} 1 - 10^{-10} & 10^{-10} \\ 10^{-10} & 1 - 10^{-10} \end{bmatrix}$$

and emission probabilities set to:

$$E = \begin{bmatrix} \theta & \epsilon \\ 1 - \theta & 1 - \epsilon \end{bmatrix}$$

Where $\epsilon = 0.001$ and $\theta = 0.01$. The most likely path was calculated using the Viterbi algorithm, and heterozygous segments 10kb or larger were retained. Heterozygous blocks within 100kb of one another in a sample strain were clustered together as a single segment to define the span of residual heterozygosity within inbred lines.

Differences in Site Frequency Spectra

If different classes of duplications have different selective impacts, we should observe clear differences in site frequency spectra, with more positively selected duplications showing fewer singleton alleles and more high frequency variants. Site frequency spectra are not normally distributed, nor can they be normalized through standard transformations, and thus require non-parametric tests. We used a two-sided Wilcoxon rank sum test to determine whether site frequency spectra were significantly different. For each comparison, we excluded tandem duplications that are present in the reference genomes as well as putative ancestral duplications, as these are likely to display biases with respect to size, propensity to capture coding sequences, and association with repetitive content. We compared site frequency spectra of the following groups within each species: duplications on the X and on the autosomes and all pairwise combinations of SNPs and duplicates on the X and autosomes. We also performed Kolmogorov-Smirnov test for comparison. In *D. simulans*, we used a χ^2 test to determine whether high frequency alleles are overrepresented among duplications on the X relative to intronic SNPs, comparing the proportion of variants as at a sample frequency of $\geq \frac{16}{17}$.

Tandem duplicates that lie in regions with residually heterozygous segments extending 1kb upstream or downstream were excluded from the SFS, resulting in unequal sample sizes for different variants. Samples with fewer than 15 strains remaining were excluded from the SFS. The SFS for intronic SNPs and for duplicates was then scaled to a sample of size 17 in *D. simulans* and 15 in *D. yakuba* according to Nielsen et al. (Nielsen et al. 2005).

Segregating Inversions

In order to check for population substructure, we aligned all SNPs in Intronic sequences from 8-30 bp which are supposed to be a neutral proxy (Halligan and Keightley 2006, Parsch et al. 2010, Clemente and Vogl 2012) and performed hierarchical clustering in R using hclust. These SNPs were intended solely to differentiate strains and were not polarized with respect to the ancestral state or otherwise filtered. We observe little evidence for population structure in *D. simulans* (Figure S12). However, we identify structure on chromosome 2 in *D. yakuba* (Figure S13), consistent with known polymorphic inversions prohibiting recombination on chromosome 2 (Lemeunier and Ashburner 1976, Llopart, Lachaise and Coyne 2005). Strains do not strictly cluster with respect to geography but rather are reticulated amongst other groups. Moreover, among duplicates we do not observe an excess of moderate frequency alleles as one would expect under population substructure given our sampling scheme (Figure S1). Thus, these strains constitute a single admixed population.

Some strains retained residual heterozygosity even after 9 generations of inbreeding, with greater residual heterozygosity in *D. yakuba* than in *D. simulans*, consistent with inversions

segregating in *D. yakuba*. These regions of residual heterozygosity can result in incorrect estimates of SFS by artificially increasing chances of observing variation. Site frequency spectra were calculated across all strais by correcting sample frequencies for ascertainment bias, excluding regions of residual heterozygosity and then projecting frequencies onto a sample size of 15 in *D. yakuba* and 17 in *D. simulans* according to (Nielsen et al. 2005). As a neutral comparison we calculated SFS for intronic SNPs (as above) and projected the SFS down to a sample size of 15 in *D. yakuba* and 17 in *D. simulans* (Figure S1).

Likelihood of shared variation through ancestry

The likelihood of shared variation through shared ancestry can be obtained through a coalescent approach. The probability that an allele does not coalesce in the time period from the present back to the speciation event that separated *D. yakuba* and *D. simulans* is $(1 - \frac{1}{2N_e})^t$. This can be approximated using $e^{\frac{-t}{2N_e}}$. We estimate θ_{pi} for putatively neutral 8-30 bp from short introns using libsequence (Thornton 2003), ignoring sites that are heterozygous and sites with missing data. For neutral intronic SNPs, $\theta_{pi} = 0.0138$ in *D. yakuba* and $\theta_{pi} = 0.0280$ in *D. simulans*. Using the mutation rate of 5.8×10^{-9} (Haag-Liautard et al. 2007), we find $N_e = (0.0138)/(4 \times 5.8 \times 10^{-9}) = 5.93 \times 10^5$ in *D. yakuba* and $N_e = (0.0280)/(4 \times 5.8 \times 10^{-9}) = 1.21 \times 10^6$ in *D. simulans*. Using $t=12\text{MY}$ (Tamura, Subramanian and Kumar 2004) and 12 generations per year, and $N_e = 1.2 \times 10^6$ from *D. simulans*, we obtain a probability of shared ancestry for an allele of 9×10^{-27} , vanishingly small. We have polarized all mutations against the putative ancestral state

using outgroup reference genomes, focusing solely on derived mutations (Rogers et al. 2014). Furthermore, the expectation of shared variation for any two alleles through shared ancestry for *D. yakuba* and *D. simulans* is expected to be low. Even large samples are not expected to harbor shared variation over such timescales (Rosenberg 2003). Thus, we expect shared variants described here to result from independent mutations, not from long standing neutral polymorphism.

Estimates of population genetic parameters by chromosome

We estimate θ_π , θ_W , and Tajima's D for all SNPs in the *D. yakuba* and *D. simulans* genomes, removing sites with missing or ambiguous data as well as heterozygous sites using libsequence (Thornton 2003). We calculate θ_π , θ_W , and Tajima's D for 5 kb windows moving in a 500 bp slide across the genome. For each window, we divide estimates by the number of sites per window with a minimum Illumina coverage depth of 20 reads, maximum coverage of 250 reads, $MQ \geq 20$, consistent with the threshold used to identify SNPs, in order to estimate θ_π and θ_W per site. We compare θ_π per site for regions surrounding derived, segregating tandem duplications with regions surrounding derived, segregating, putatively neutral intronic SNPs from 8-30 bp of short introns, excluding windows with less than 4000 bp out 5000 bp that could be assayed for SNPs. We exclude second SNPs in a single 5 kb window, and exclude SNPs and duplicates that are found in the centromeric regions, which have unusually low diversity (Figure S2-S11). We compared diversity at the locus of SNPs to diversity at the locus of duplication by chromosome with a Wilcox rank sum test (Table S4) and estimated

mean diversity from 50 kb upstream to 50kb downstream of a mutant (Figure 2-3). Tests of nucleotide diversity exclude a cluster of multiple duplications from 8.45 Mb-8.55 Mb which has abnormally low diversity (Figure S9).

Bibliography

- Clemente F, Vogl C. 2012. Unconstrained evolution in short introns? - an analysis of genome-wide polymorphism and divergence data from *Drosophila*. J. Evol. Biol. 25:1975–1990.
- Emerson JJ, Cardoso-Moreira M, Borevitz JO, Long M. 2008. Natural selection shapes genome-wide patterns of copy-number polymorphism in *Drosophila melanogaster*. Science. 320:1629–1631.
- Haag-Liautard C, Dorris M, Maside X, Macaskill S, Halligan DL, Houle D, Charlesworth B, Keightley PD. 2007. Direct estimation of per nucleotide and genomic deleterious mutation rates in *Drosophila*. Nature. 445:82–85.
- Halligan DL, Keightley PD. 2006. Ubiquitous selective constraints in the *Drosophila* genome revealed by a genome-wide interspecies comparison. Genome Res. 16:875–884.
- Lemeunier F, Ashburner MA. 1976. Relationships within the melanogaster species subgroup of the genus *Drosophila* (Sophophora). II. Phylogenetic relationships between six species based upon polytene chromosome banding sequences. Proc. R. Soc. Lond., B, Biol. Sci. 193:275–294.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 25:2078–2079.
- Llopart A, Lachaise D, Coyne JA. 2005. Multilocus analysis of introgression between two sympatric sister species of *Drosophila*: *Drosophila yakuba* and *D. santomea*. Genetics. 171:197–210.
- McKenna A, Hanna M, Banks E, et al. (11 co-authors). 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res. 20:1297–1303.
- Nielsen R, Bustamante C, Clark AG, et al. (13 co-authors). 2005. A scan for positively selected genes in the genomes of humans and chimpanzees. PLoS Biol. 3:e170.

- Parsch J, Novozhilov S, Saminadin-Peter SS, Wong KM, Andolfatto P. 2010. On the utility of short intron sequences as a reference for the detection of positive and negative selection in *Drosophila*. *Mol. Biol. Evol.* 27:1226–1234.
- Rogers RL, Bedford T, Lyons AM, Hartl DL. 2010. Adaptive impact of the chimeric gene *Quetzalcoat1* in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences, USA*. 107:10943–10948.
- Rogers RL, Cridland JM, Shao L, Hu TT, Andolfatto P, Thornton KR. 2014. Landscape of standing variation for tandem duplications in *drosophila yakuba* and *drosophila simulans*. *arXiv preprint arXiv:1401.7371*. .
- Rogers RL, Hartl DL. 2012. Chimeric genes as a source of rapid evolution in *Drosophila melanogaster*. *Mol. Biol. Evol.* 29:517–529.
- Rosenberg NA. 2003. The shapes of neutral gene genealogies in two species: probabilities of monophyly, paraphyly, and polyphyly in a coalescent model. *Evolution*. 57:1465–1477.
- Tamura K, Subramanian S, Kumar S. 2004. Temporal patterns of fruit fly (*Drosophila*) evolution revealed by mutation clocks. *Mol. Biol. Evol.* 21:36–44.
- Thornton K. 2003. libsequence: a C++ class library for evolutionary genetic analysis. *Bioinformatics*. 19:2325–2327.

Table S1: Number of duplicated regions detected in *D. yakuba* and *D. simulans*

	<i>D. yakuba</i>	<i>D. simulans</i>
Whole gene	248	296
Partial gene	745	462
Intergenic	745	577

Table S2: Wilcoxon Rank Sum Tests of Site Frequency Spectra

Species	Type	Type	W	Adjusted P -value
<i>D. yakuba</i>	Autosomal SNPs	Autosomal Duplicates	212	$3.507 \times 10^{-6**}$
	X-linked SNPs	X-linked Duplicates	211	$4.781 \times 10^{-4**}$
	Autosomal Duplicates	X-linked Duplicates	172	0.0128*
<i>D. simulans</i>	Autosomal SNPs	Autosomal Duplicates	268	$2.981 \times 10^{-6**}$
	X-linked SNPs	X-linked Duplicates	113	0.2897
	Autosomal Duplicates	X-linked Duplicates	183.5	0.1848

* $P < 0.05$, ** $P < 0.01$

SNPs are derived from 8-30 bp of short first introns ≤ 100 bp.

Table S3: Kolmogorov-Smirnov Tests of Site Frequency Spectra

Species	Type	Type	D	Adjusted P -value
<i>D. yakuba</i>	Autosomal SNPs	Autosomal Duplicates	0.9333	$3.868 \times 10^{-7**}$
	X-linked SNPs	X-linked Duplicates	0.800	$5.235 \times 10^{-5**}$
	Autosomal Duplicates	X-linked Duplicates	0.5333	0.02625*
<i>D. simulans</i>	Autosomal SNPs	Autosomal Duplicates	0.8824	$4.808 \times 10^{-7**}$
	X-linked SNPs	X-linked Duplicates	0.2941	0.4654
	Autosomal Duplicates	X-linked Duplicates	0.3529	0.2402

* $P < 0.05$, ** $P < 0.01$

SNPs are derived from 8-30 bp of short first introns ≤ 100 bp.

Table S4: Reduced Diversity Surrounding Tandem Duplications

Species	Chrom	W	single tailed P
<i>D. yakuba</i>	2L	27768	0.9993
	2R	27768	0.9888
	3L	17354.5	0.4797
	3R	76908.5	0.0064
	X	16660	0.2674
<i>D. simulans</i>	2L	31594	1.259×10^{-4}
	2R	56665	0.453
	3L	41054.5	0.01074
	3R	42493	3.166×10^{-6}
	X	14603.5	0.01815

Table S5: Gene ontology categories overrepresented in both *D. yakuba* and *D. simulans*

Functional Category	<i>D. yakuba</i> EASE	<i>D. simulans</i> EASE
Chitin metabolism or cuticle	2.00	0.97
Immune response	1.44	1.59
Drug and toxin metabolism	1.37	2.32
Chemosensation	1.12	1.37
Multiple Independent Duplications		
Chorion and oogenesis	1.79	1.84
Sensory processing	1.23	1.41
Immune response	1.11	3.35
High Frequency Duplicates		
Endopeptidases	-	-

Table S6: Estimated Number of Segregating Duplications on X and major Autosomes

	<i>D. yakuba</i>	<i>D. simulans</i>
Genome wide θ_W for tandem duplications	383	264
$E[S]$	5700	3800
σ_S	497	344
$E[S] + 2\sigma_S$	6800	4500
Genomic Coverage	13.4%	9.7%

Table S7: Number of duplications necessary to cover segments of the genome

Percent Covered	Lower Bound (95% CI)	Upper Bound (95%CI)
5%	2,358	2,660
10%	4,912	5,360
25%	13,668	14,410
50%	33,191	34,427
90%	119,799	113,767

Table S8: Multiply duplicated genes (≥ 4 duplications)

Species	Gene	Number
<i>D. yakuba</i>	GE10684-PA	4
	GE13282-PA	4
	GE18810-PA	4
	GE18813-PA	4
	GE18814-PA	4
	GE20773-PA	4
	GE20774-PA	4
	GE25839-PA	4
	GE18811-PA	6
	GE18812-PA	6
<i>D. simulans</i>	CG2174-PD	4
	CG33466-PA	4
	CG4250-PA	4
	CG42566-PA	4
	CG33162-PA	15
	CG32022-PA	16
	CG5939-PA	27
	CG6533-PA	29
	CG6511-PA	32
	CG6517-PA	32
	CG6519-PA	32
	CG6524-PA	32

Supplementary Figures




Figure S1: Site frequency spectra for SNPs (black) and tandem duplications (grey) on the A) X and B) autosomes in *D. yakuba* and on the C) X and D) autosomes in *D. simulans*. Tandem duplications in *D. yakuba* and on the *D. simulans* autosomes show an excess of low frequency variants, consistent with detrimental phenotypic effects. The *D. simulans* X shows an excess of high frequency variants, consistent with widespread selection favoring duplicates on the X.




Figure S2: Nucleotide Diversity θ_π and Tajima's D chromosome 2L in *D. yakuba* for 5kb windows with a 500 bp slide, showing only windows with 1 kb or more of sequence with coverage sufficient to call SNPS. Tajima's D is negatively skewed, consistent with recent population expansion and mean diversity θ_π is low surrounding centromeric regions. Locations and diversity values for duplications are marked in red.




Figure S3: Nucleotide Diversity θ_π and Tajima's D chromosome 2R in *D. yakuba* for 5kb windows with a 500 bp slide, showing only windows with 1 kb or more of sequence with coverage sufficient to call SNPS. Tajima's D is negatively skewed, consistent with recent population expansion and mean diversity θ_π is low surrounding centromeric regions. Locations and diversity values for duplications are marked in red.




Figure S4: Nucleotide Diversity θ_π and Tajima's D chromosome 3L in *D. yakuba* for 5kb windows with a 500 bp slide, showing only windows with 1 kb or more of sequence with coverage sufficient to call SNPS. Tajima's D is negatively skewed, consistent with recent population expansion and mean diversity θ_π is low surrounding centromeric regions. Locations and diversity values for duplications are marked in red.




Figure S5: Nucleotide Diversity θ_π and Tajima's D chromosome 3R in *D. yakuba* for 5kb windows with a 500 bp slide, showing only windows with 1 kb or more of sequence with coverage sufficient to call SNPS. Tajima's D is negatively skewed, consistent with recent population expansion and mean diversity θ_π is low surrounding centromeric regions. Locations and diversity values for duplications are marked in red.




Figure S6: Nucleotide Diversity θ_π and Tajima's D for the X chromosome in *D. yakuba* for 5kb windows with a 500 bp slide, showing only windows with 1 kb or more of sequence with coverage sufficient to call SNPS. Tajima's D is negatively skewed, consistent with recent population expansion and mean diversity θ_π is low surrounding centromeric regions. Locations and diversity values for duplications are marked in red.




Figure S7: Nucleotide Diversity θ_π and Tajima's D chromosome 2L in *D. simulans* for 5kb windows with a 500 bp slide, showing only windows with 1 kb or more of sequence with coverage sufficient to call SNPS. Tajima's D is negatively skewed, consistent with recent population expansion and mean diversity θ_π is low surrounding centromeric regions. Locations and diversity values for duplications are marked in red.




Figure S8: Nucleotide Diversity θ_π and Tajima's D chromosome 2R in *D. simulans* for 5kb windows with a 500 bp slide, showing only windows with 1 kb or more of sequence with coverage sufficient to call SNPS. Tajima's D is negatively skewed, consistent with recent population expansion and mean diversity θ_π is low surrounding centromeric regions. Locations and diversity values for duplications are marked in red.




Figure S9: Nucleotide Diversity θ_π and Tajima's D chromosome 3L in *D. simulans* for 5kb windows with a 500 bp slide, showing only windows with 1 kb or more of sequence with coverage sufficient to call SNPS. Tajima's D is negatively skewed, consistent with recent population expansion and mean diversity θ_π is low surrounding centromeric regions. Locations and diversity values for duplications are marked in red. Chromosome 3L is strongly affected by a cluster of duplications over 30 independent duplications of a genomic segment at roughly 8.5Mb which shows greatly reduced diversity.




Figure S10: Nucleotide Diversity θ_π and Tajima's D chromosome 3R in *D. simulans* for 5kb windows with a 500 bp slide, showing only windows with 1 kb or more of sequence with coverage sufficient to call SNPS. Tajima's D is negatively skewed, consistent with recent population expansion and mean diversity θ_π is low surrounding centromeric regions. Locations and diversity values for duplications are marked in red.




Figure S11: Nucleotide Diversity θ_π and Tajima's D for the X chromosome in *D. simulans* for 5kb windows with a 500 bp slide, showing only windows with 1 kb or more of sequence with coverage sufficient to call SNPS. Tajima's D is negatively skewed, consistent with recent population expansion and mean diversity θ_π is low surrounding centromeric regions. Locations and diversity values for duplications are marked in red.




Figure S12: Hierarchical clustering of intronic SNP data for *D. simulans* shows little population structure.




Figure S13: Hierarchical clustering of intronic SNP data for *D. yakuba* shows population structure on chromosome 2L (B) and 2R (C), consistent with known inversions segregating on chromosome 2. Samples do not cluster strictly with respect to geography (A,D-E), indicating widespread gene flow between geographic locations and a single admixed population.